

The distances of short-hard GRBs and the SGR connection

Ehud Nakar¹, Avishay Gal-Yam^{1,2}, Tsvi Piran^{1,3} and Derek B. Fox¹

1. *Division of Physics, Mathematics and Astronomy, Caltech, Pasadena, 91125, USA*
2. *Hubble fellow*
3. *Racah Institute for Physics, The Hebrew University, Jerusalem, 91904, ISRAEL*

ABSTRACT

We present a search for nearby ($D \lesssim 100$ Mpc) galaxies in the error boxes of six well-localized short-hard gamma-ray bursts (GRBs). None of the six error boxes reveals the presence of a plausible nearby host galaxy. This allows us to set lower limits on the distances and, hence, the isotropic-equivalent energy of these GRBs. Our lower limits are around 1×10^{49} erg (at 2σ confidence level); as a consequence, some of the short-hard GRBs we examine would have been detected by BATSE out to distances greater than 1 Gpc and therefore constitute a bona fide cosmological population. Our search is partially motivated by the December 27, 2004 hypergiant flare from SGR 1806–20, and the intriguing possibility that short-hard GRBs are extragalactic events of a similar nature. Such events would be detectable with BATSE to a distance of 50 Mpc, and their detection rate should be comparable to the actual BATSE detection rate of short-hard GRBs. The failure of our search, by contrast, suggests that such flares constitute less than 15% of the short-hard GRBs ($< 40\%$ at 95%-confidence). We discuss possible resolutions of this discrepancy.

Subject headings: Gamma-Ray: Bursts

1. Introduction

After a decade of rapid progress in our understanding of gamma-ray bursts (GRBs; see Piran 2004 for a recent review), the origin of short-hard GRBs is still a mystery. The short-hard GRB class (Kouveliotou et al. 1993) makes up a quarter of the entire GRB population observed by BATSE¹ with an all-sky detection rate of $\approx 170 \text{ y}^{-1}$ (Meegan et al., 1997). The revolution in our understanding of the long-duration GRBs, triggered by

¹<http://www.batse.msfc.nasa.gov/batse/>

the detection of their long-lived counterpart (afterglow) emission, has so far passed over the short-hard GRBs, lacking similar afterglow detections². With no afterglow detections, we have only indirect clues about the typical distance to short-hard GRBs: the almost-isotropic sky distribution (Briggs et al. 1996; Balazs et al. 1998; Magliocchetti et al. 2003) and small value of $\langle V/V_{max} \rangle$ (Katz & Canel 1996; Schmidt 2001; Guetta & Piran 2004) for the population suggests a cosmological origin. Even so, the distance and energy scale of the events remains highly uncertain.

Soft gamma-ray repeaters (SGRs) are compact sources of persistent X-ray emission and repeating bursts of soft gamma-rays (see Woods & Thompson 2004 for a review). They are believed to be highly-magnetized young neutron stars (known as magnetars; Duncan & Thompson 1992; Paczyński 1992; Katz 1994; Thompson & Duncan 1995, see also Katz 1982 for related early ideas). At infrequent intervals, these sources emit powerful flares of high energy radiation, much larger than their usual bursts, releasing more than 10^{44} ergs in the form of gamma-rays alone. These “giant flares” are characterized by a short ($\lesssim 0.5$ sec) and very intense burst that is followed by a long (~ 100 sec) pulsating soft tail. During ~ 30 years of observations, two of the four confirmed SGRs (SGR 0526–66 and SGR 1900+14) produced a single giant flare each.

On December 27, 2004 a third source, SGR 1806–20, erupted in a giant flare (Borkowski et al. 2004; Mazets et al. 2005; Hurley et al. 2005; Palmer et al. 2005). The initial pulse fluence was about 1 erg/cm^2 . At a distance of 15 kpc (Corbel & Eikenberry 2004; see however Figer et al. 2004; Cameron et al. 2005; § 5) this fluence corresponds to a total energy output of 3×10^{46} erg, about a hundred times the energy of the two previous giant flares, making this a hypergiant flare (hereafter HGF). As seen in previous giant flares this event showed a short and intense pulse followed by a long pulsating soft tail.

Duncan (2001) pointed out that the initial hard pulse of an extragalactic giant flare would be detected by BATSE as a short-hard GRB. He considered flares with a luminosity similar to that of the giant flare of SGR 0526–66, finding that they would be detected to a distance of ~ 10 Mpc and could account for only a small fraction of the short GRBs observed by BATSE. Eichler (2002) suggested that “total field reversal” in a magnetar could produce an HGF with an energy output of $\sim 10^{46}$ to 10^{47} erg in a catastrophic event involving the near-total destruction of the magnetic dipole moment of the star. Such an HGF would be detected as a short-hard GRB out to a distance of ~ 100 Mpc. However, according to Eichler’s model, an HGF consumes most of the exterior magnetic energy of the magnetar and

²The likely X-ray afterglow of the short-hard burst GRB 050509b was discovered by *SWIFT* (Gehrels et al. 2005) while this paper was being refereed. We briefly discuss the implications of this event in § 5.

can occur only once or twice during the SGR’s lifetime, implying an HGF rate of 10^{-3} y^{-1} per Milky Way-size Galaxy. Eichler (2002) therefore also concluded that HGFs would constitute only a small fraction of the short GRBs in the BATSE sample.

The observation of an HGF with an energy larger than 10^{46} erg after only ~ 30 years of observations in our Galaxy has forced a revision of this picture. Boggs et al. (2005) reiterated the possible connection between HGFs and short GRBs. Dar (2005) pointed out that an HGF rate of $\sim 3 \times 10^{-2} \text{ y}^{-1}$ in the Milky Way implies an observed all-sky rate of roughly one short GRB per day at BATSE sensitivity. This rate is comparable to the observed rate of short GRBs, suggesting that all short GRBs may be HGFs from SGRs.

In this letter we report on a search for nearby star-forming host galaxies in the error boxes of six well-localized short-hard GRBs. We use our limits on the brightness of any such host galaxies to derive minimal distances and isotropic-equivalent energy outputs for these bursts. We use the measured flux of the Dec. 27 HGF in the BATSE detection window (50–300 keV), verified by our own independent estimation derived using the observed backscatter of HGF X-rays from the moon (Golenetskii et al. 2004), to estimate the BATSE detection rate of extragalactic HGFs and to compare this prediction with the results of our search.

In the sections below, we estimate the energy output of the recent HGF from SGR 1806-20 in the BATSE detection window (§2), calculate the expected BATSE detection rate of extragalactic HGFs (§3), and describe our search for nearby host galaxies of short-hard GRBs (§4). We conclude with a discussion of the implications of our work for theories of the short-hard GRBs and, in particular, their relation to the SGRs (§5). A Hubble constant of $H_0 = 70 \text{ km/sec/Mpc}$ is assumed throughout the paper.

2. The fluence of the SGR 1806–20 hypergiant flare

The Coranas-F spacecraft ³, a solar activity observatory in a low-earth orbit, was occulted by the Earth at the time of the flare. The moon, at this time, was full with respect to Coranas-F and the flare. This coincidence enabled Helicon, a γ -ray spectrometer on Coranas-F, to measure the backscatter of the flare from the moon without being saturated. Helicon measured a total backscattered fluence, $f_{bs} = 7.5 \times 10^{-7} \text{ erg/cm}^2$ in the 25 – 400 keV range (Golenetskii et al. 2004).

The spectrum of the backscattered radiation, a broad hump around 100 keV, is consistent with the spectrum of reflected X-rays and gamma-rays from cold matter (Lightman &

³<http://coronas.izmiran.rssi.ru/F/>

White 1988). The reflection of low energy photons below ~ 50 keV is suppressed by atomic absorption (Lightman & White 1988; George & Fabian 1991), while above this energy the dominant process is electron scattering. In this regime, reflection of photons of increasing energy ($E_\gamma \gtrsim 300$ keV) is progressively more suppressed by the sharply decreasing Klein-Nishina cross section for backscattering and by the photon energy loss at each scatter. The albedo for monochromatic radiation with a normal incidence angle to a semi infinite plane parallel slab is $\approx 35\%$ for 100 keV photons and $\approx 20\%$ for 300 keV photons, while it drops to a few percent at 1 MeV (Imamura & Epstein 1987). In all these cases the reflected spectrum peaks below 300 keV due to the Klein-Nishina energy loss at each scatter. The sharp drop of the albedo above 300 keV implies that unless the spectrum is very hard, the emerging fluence in the energy window of Helicon closely reflects the incoming fluence in a similar window.

Assuming a flat spectrum ($F_\nu \propto \text{const}$), we approximate the backscatter albedo of the moon in Helicon’s window as $a \approx 0.25$ (i.e. the ratio between the incoming fluence and the reflected fluence in the energy window of Helicon). Unless the spectrum is extremely hard and all the energy is emitted above 0.5 MeV the exact incident spectral shape and the exact geometry of the scattering should introduce a correction factor of order unity to this value. Now we can approximate the fluence of the HGF as: $S_{25-400} = 8S_{bs}/(A_m^2 a) \approx 0.3 \text{ erg/cm}^2$ where $A_m = 0.009 \text{ rad}$ is the angular size of the moon from earth. This fluence implies, using a distance of 15 kpc, an isotropic equivalent energy of $E_{25-400} \approx 0.8 \times 10^{46} \text{ erg}$.⁴

3. The detection rate of extragalactic hypergiant flares by BATSE

As young neutron stars, SGRs are remnants of recent core-collapse SNe. Thus, their abundance should be proportional to the star formation rate (SFR). Using the SFR in the local universe, $SFR_{loc} \approx 0.02 M_\odot \text{ y}^{-1} \text{ Mpc}^{-3}$ (Pérez-González et al. 2003; Schiminovich et al. 2004), the rate of HGFs in the local universe can be estimated as:

$$\dot{N}_{HGF} = 2 \times 10^{-4} \text{ y}^{-1} \text{ Mpc}^{-3} \frac{\dot{N}_{HGF,MW}}{1/30 \text{ y}^{-1}} \frac{SFR_{loc}}{0.02 M_\odot \text{ y}^{-1} \text{ Mpc}^{-3}} \frac{3 M_\odot \text{ y}^{-1}}{SFR_{MW}}, \quad (1)$$

where SFR_{MW} is the SFR in the Milky Way ($3_{-2}^{+6} M_\odot \text{ y}^{-1}$; Timmes et al. 1997; Miller & Scalo 1979). BATSE’s effective threshold⁵ for triggering on a time scale of 256 ms is an

⁴The measurements of fluence and spectrum of the flare was published (Hurley et al. 2005, Palmer et al. 2005) while this paper was in the refereeing process. Our estimate of the fluence in the 25 – 400 keV range is within the errors of these measurements.

⁵http://coss.c.gsfc.nasa.gov/batse/BATSE_Ctlg/trig_sen.html

energy flux of 10^{-7} erg/cm²/sec in the 50 – 300 keV energy band⁶ Since the duration of the HGF initial pulse was shorter than 0.25 sec, this threshold corresponds to a fluence of $S_{lim,50-300} \approx 2.5 \times 10^{-8}$ erg/cm². Fortunately, the albedo of the moon peaks around BATSE’s detection window, enabling us to estimate the fluence of the HGF at this window, $S_{50-300} \approx 0.3$ erg/cm² (§2). This implies a maximal distance for detection by BATSE of ≈ 50 Mpc. Thus, the all-sky rate of similar extragalactic HGFs above BATSE’s detection threshold is:

$$\dot{N}_{HGF}^{obs} \approx 130 \text{ y}^{-1} \left(\frac{R_{SGR}}{15 \text{ kpc}} \right)^3 \left(\frac{S_{50-300}/0.3}{S_{lim,50-300}/2.5 \times 10^{-8}} \right)^{\frac{3}{2}} \frac{\dot{N}_{HGF,MW}}{1/30 \text{ y}^{-1}} \frac{SFR_{loc}}{0.02 M_{\odot} \text{ y}^{-1} \text{ Mpc}^{-3}} \frac{3 M_{\odot} \text{ y}^{-1}}{SFR_{MW}}, \quad (2)$$

where R_{SGR} is the distance to SGR 1806-20. This result is insensitive to the introduction of beaming (Dar 2005) since in this case $\dot{N}_{HGF,MW}$ is the rate of HGFs in the Milky Way which are pointing towards us. This rate is comparable to the all-sky rate of short GRBs in BATSE’s sample, $\dot{N}_{SGRB}^{obs} \approx 170 \text{ y}^{-1}$, suggesting that *all* short GRBs are extragalactic HGFs.

4. A search for nearby host galaxies associated with short-hard GRBs

We have compiled a list of all IPN short-hard GRBs with fine localizations (3σ error box area ≤ 100 arcmin²) which are also at high Galactic latitude ($\delta > 20$) to avoid the complications caused by extinction and source confusion in Galactic fields. We have found six bursts which satisfy our criteria⁷.

We first consider the possibility that short-hard GRBs are physically associated with regions of star formation, as are long-duration GRBs, soft gamma-ray repeaters (SGRs), and progenitors proposed by several alternate models. In this framework we obtain a limit on the distance of a putative host galaxy using the proportionality of the SFR to the UV luminosity of galaxies in the nearby universe (Kennicutt 1998). Recent results from the Galaxy Evolu-

⁶This threshold assumes a mildly hard spectrum - a power law with a photon index of -1.5. The thermal spectrum of the flare, which is approximately flat in BATSE’s window, should introduce a correction of order unity.

⁷Note that one of these bursts (GRB 020603) has a duration of 1.5 sec in the 25 – 100 keV band as observed by Ulysses (Hurley et al 2002). Since this energy band is similar to the two lowest BATSE bands and since the duration of GRBs is typically shorter at higher energy bands, we consider this burst as a viable short burst. Our results remain unchanged if we exclude this burst from the sample.

tion Explorer (GALEX⁸ show that the UV luminosity function of nearby galaxies (Wyder et al. 2005) is such that 68%[95%] of the UV luminosity (and thus, SFR) is associated with galaxies brighter than $M_{UV} = -17[-14]$ (which translates to an SFR of $0.33[0.02]M_{\odot}$ per year). Thus the hypotheses that a galaxy fainter than $M_{UV} = -17[-14]$ is the host of a short GRB can be rejected at $1\sigma[2\sigma]$. Given these limits on the UV brightness of possible hosts, the *observed* luminosities of galaxies within GRB error boxes impose lower limits on their distances. If nearby, these galaxies would be less luminous than the limit defined above, and thus unlikely to host a GRB. The weakest limit is obtained for the brightest galaxy in the field and we conservatively adopt this as the lower limit on the distance to each GRB, from which we derive a lower limit on its energy.

Where possible, we use available UV imaging from the GALEX mission to find the SFR of putative host galaxies. In those cases where UV data are not available, we derive the SFR of each putative host galaxy from its B band magnitude and its $U-V$ color (Kennicutt 1998). We verify that this method produces conservative results when compared to the GALEX UV analysis. Finally, we briefly consider also the possibility that short-hard GRBs follow the host galaxies B -band or R -band luminosity (rather than the UV/SFR).

GALEX UV data were available for three of the six bursts in our sample. We have extracted NUV magnitudes and errors for all the sources detected by GALEX within these error boxes from the Multy-Mission Science Archive⁹ at STScI. We then translated the NUV flux to SFR (Kenicutt 1998) and deduced the minimal distance to the GRB by requiring that each galaxy has an SFR above the threshold described above. In Table 1 we give the minimal distances derived from the brightest GALEX source in each error box.

For the four localizations for which GALEX data are not available, we have extracted XDSS POSS II plates¹⁰. We have also carried out plate-based analysis for two GRBs for which GALEX data are available, to check for consistency. Red plates were available for all fields, and for most of them we have also examined blue and infrared plates (Table 1). We located the brightest galaxy within the boundaries of the error box (see Fig. 1 for an example). In all cases, the brightest galaxy was the same one in all bands. Additional available data indicated that several of the nominally brightest galaxies were unlikely to be nearby host galaxies. In those cases we have considered instead either the next brightest galaxy in the field or if none was found a hypothetical galaxy at the plate detection limit. See Table 1 for details.

⁸<http://www.galex.caltech.edu/>

⁹<http://archive.stsci.edu/>

¹⁰<http://salish.dao.nrc.ca:8080/dss/>

Using the XDSS astrometry, we then retrieved the cataloged B and V magnitudes of the galaxies of interest, using the Naval Observatory Merged Astrometric Data set (NOMAD; Zacharias et al. 2004) and USNO-B1 (Monet et al. 2003) catalogs¹¹. When available, we have also extracted g , r and i magnitudes from the DPOSS survey¹² (Djorgovski et al. 2003) and converted these to B and V magnitudes using transformations from Smith et al. (2002). For two bursts (021201 and 020603), Henden (2002a, 2002b) provides photometric calibrations of the field. From these, we retrieve accurate B and V magnitudes of the relevant galaxies. The uncertainty in $B - V$ from Henden is $\delta(B - V) = 0.05$ and $\delta(B - V) = 0.005$ for the brightest galaxies in the fields of GRB 021201 and 020603, respectively. The uncertainty in DPOSS-based $B - V$ values we estimate as $\delta(B - V) \sim 0.1$, including both zero point errors and the uncertainty introduced by the Smith et al. (2002) transformation. For one burst for which our $B - V$ color is based solely on NOMAD cataloged values (031214), we set a conservative error of $\delta(B - V) \sim 0.2$, equal to the maximum offset we find when comparing NOMAD values to either DPOSS or Henden calibration available for the four other bursts. We list our adopted B and V magnitudes and the uncertainty in $B - V$ for the brightest galaxy detected in each GRB field in Table 1.

Using the correlation derived by Gavazzi, Boselli and Kennicutt (1991; see their Fig. 1) we use the $B - V$ colors of these galaxies to derive their U -band magnitude and thus their $U - V$ color. With the $U - V$ color in hand, we use the results of Kennicutt (1998; see Fig. 2) to derive the star formation rate in each galaxy, in solar masses per year per 10^{10} solar luminosities in the B -band. The $B - V$ uncertainty is properly propagated. Finally, we use this to find the distance at which the B -band luminosity of each putative host galaxy (derived from the apparent B magnitude and the distance) is equivalent, through the Kennicutt relation, to an SFR of $0.33[0.02]M_{\odot}$ per year (our SFR threshold distances appear in table 2).

To verify the consistency of the SFRs derived from optical colors using the Kenicutt relation with those derived directly from GALEX UV data, we conducted both types of analyses for two bursts and compared the resulting SFR limits. We find that our optical analysis is conservative, in the sense that the SFR value derived for the brightest galaxy in each field from optical observations is higher than the values we get from GALEX UV observations. We are thus reassured that our analysis of the bursts without GALEX data gives conservative lower limits on the distance and energy of the relevant GRBs. As another test of the analysis we have measured the redshift, $z = 0.14$, of the brightest galaxy in the

¹¹<http://www.nofs.navy.mil/nomad/>

¹²<http://dposs.ncsa.uiuc.edu>

field of GRB 000607 (Fig. 1). The derived distance, 580 Mpc, is significantly larger than the minimal distance that we have estimated.

These minimal distance to putative host galaxies result in lower limits for the isotropic-equivalent energy output of each burst, $10^{48} - 10^{49}$ erg at 2σ and $10^{49} - 10^{50}$ at 1σ (with confidence levels given for each individual burst). In table 2 we list these distances and the corresponding lower limits on the bolometric luminosity (when available) and on the luminosity in the BATSE window for triggering (50–300 keV). The lower limits we obtain show that none of the six short bursts we have examined were produced by an event that is similar to the recent HGF from SGR 1806–20.

We have repeated the same analysis (to all the sample except for GRB 790613) assuming that the progenitors of short-hard GRBs follow the blue light or that they follow the red light. Namely, we found the minimal distance to any putative galaxy assuming that its absolute magnitude is $0.33[0.02]L_*$ in the blue band (using the *B*-band luminosity function from the 2dF Galaxy Redshift Survey; Croton et al. 2005) and in the red band (using the *R*-band luminosity function from the Sloan Digital Sky Survey; Blanton et al. 2003). The results in both cases are qualitatively similar to the limits we obtain under the assumption that the short-hard GRBs follow the SFR.

5. Discussion

We have obtained lower bounds on the isotropic equivalent energy output of six short-hard GRBs under the assumption that their rate follows the UV luminosity (star formation rate), blue luminosity (as does the rate of type Ia SNe; Tammann 1970, van den Bergh & Tammann 1991) or red luminosity of their host galaxies. Under all assumptions we get comparable limits, suggesting that our results generally hold in any case that short-hard GRBs are spatially associated with host galaxies and follow their light¹³. With a minimal energy of $10^{48} - 10^{49}$ erg in the detection window of BATSE (Table 1) these bursts are detectable to distances larger than ~ 1 Gpc. Therefore these bursts are part of a cosmological population of short-hard GRBs, in agreement with the low $\langle V/V_{max} \rangle$ value of the short-hard GRB population (Katz & Canel 1996; Schmidt 2001; Guetta & Piran 2004).

¹³Note that this is not the case in the NS-NS merger scenario where bursts are not necessarily associated with nearby galaxies. In this scenario the coalescence can occur at a distance of ~ 200 kpc from the host galaxy (Narayan et al. 1992). Therefore, for bursts that are closer than 10 Mpc, the host galaxies may be outside the error boxes we examined. However, in the NS-NS merger scenario bursts are expected to be detectable to much larger distances.

With a bolometric energy output larger than 10^{49} erg over a fraction of a second, their peak luminosities are comparable to those of long duration GRBs. These results suggest that the mechanism that produces the majority of short GRBs is related to the mechanism that produces long GRBs, as is also suggested by the similarity of the temporal (Nakar & Piran 2002) and spectral (Ghirlanda et al. 2004) structure of short-hard GRBs and the first two seconds of long GRBs.

The expected detection rate of extragalactic HGFs similar to the December 27, 2004 event as short-hard bursts by BATSE (Eq. 2) is comparable to the total rate of such GRBs. On the other hand, the lower limits we obtain for our sample of short bursts are at least two orders of magnitude above the energy emitted during the December 27 event. Thus, our results suggest that less than 15% of BATSE short-hard GRBs are December 27-type HGFs, and excludes at 95% confidence level the possibility that more than 40% of the short GRBs are HGFs.

These findings together leave us with four options, none compelling. First, it is possible that the short-hard GRB population is in fact composed of two unrelated phenomena, one local and one cosmological. By coincidence those two phenomena have comparable detection rates at the BATSE threshold as well as similar temporal and spectral properties. In this scenario the local population would preferentially enrich the sample close to and below the BATSE threshold. This scenario is only marginally consistent with our null detection of nearby star-forming galaxies associated with six short GRBs.

A second possibility is that all short GRBs are HGFs but that some HGFs emit as much as 10^{49} erg of high-energy radiation. In this scenario the luminosity function of HGFs should decrease rapidly with energy so that the observed number of HGFs with $E = 10^{46}$ erg would be comparable to that of HGFs with $E = 10^{49}$ erg despite their being visible out to very different distances. A disturbing question in this scenario is how do isolated neutron stars produce such gigantic bursts.

A third possibility is that we have observed an unlikely event. That is, even though the true rate of HGFs is one per SGR lifetime (one per ~ 1000 years), we have seen one after only 30 years of observations. This implies that HGFs are unique events in the life of SGRs, as predicted by Eichler (2002), making HGFs a small fraction of short-hard GRBs.

Finally, a fourth alternative is that the distance to SGR 1806–20 has been overestimated, and is actually about 6 kpc. The distance determination of this SGR is controversial, and based on its association with a star cluster. Corbel & Eikenberry (2004) find it to be at 15 kpc while Figer et al. (2004) suggest that it is at a distance of 11.8 kpc, with the only robust evidence as to its distance being but a lower limit of 6.4 kpc (Cameron et al. 2005).

At a smaller distance, the December 27 event is demoted to the rank of bright giant flare. With energy lower by an order of magnitude compared to the current estimate, the detection rate of similar extragalactic flares would decrease by a factor of ~ 30 , making them a small fraction of short GRBs.

While this paper was being refereed, *SWIFT* has localized the short-hard burst GRB 050509b (Geherles et al. 2005). While the identity of the host galaxy of this burst has not been conclusively determined (e.g., Bloom et al 2005, Fox et al 2005) it is clearly at or above $z = 0.22$ ($D=900$ Mpc). Thus, this event strengthens our results, suggesting that most short-hard bursts are at cosmological distances. Furthermore, the low localization rate of short-hard bursts achieved by *SWIFT* so far ($\lesssim 4 \text{ yr}^{-1}$) suggests that the sample size of well-localized short-hard bursts is not likely to rapidly increase. Statistical studies of host galaxies will therefore probably continue to rely heavily on “historical” short-hard burst samples, such as the one we have analyzed here.

We would like to thank E. S. Phinney, R. Sari, M. Kamionkowski, S. Furlanetto, J. Katz, A. Pickles and S. Eikenberry, for helpful discussions and remarks. We thank J.L. Atteia for directing our attention to GRB 790613. A.G. acknowledges support by NASA through Hubble Fellowship grant #HST-HF-01158.01-A awarded by STScI, which is operated by AURA, Inc., for NASA, under contract NAS 5-26555. This research was partially supported by a grant from the US-Israel BSF.

REFERENCES

- Balazs, L. G., Meszaros, A., & Horvath, I. 1998, *A&A*, 339, 1
- Barat, C., et al. 1984, *ApJ*, 280, 150
- Blanton, M. R., et al. 2003, *ApJ*, 592, 819
- Bloom, J. S., et al. 2005, *ArXiv Astrophysics e-prints*, arXiv:astro-ph/0505480
- Briggs, M. S., et al. 1996, *ApJ*, 459, 40
- Cameron, P. B., et al. 2005, *Nature*, 434, 1112
- Corbel, S., & Eikenberry, S. S. 2004, *A&A*, 419, 191
- Croton, D. J., et al. 2005, *MNRAS*, 356, 1155
- Dar, A., 2005, *GCN circ.* 2942

- Djorgovski, S. G., Carvalho, R. R., Gal, R. R., Odewahn, S. C., Mahabal, A. A., Brunner, R., Lopes, P. A. A., & Kohl Moreira, J. L. 2003, *Bulletin of the Astronomical Society of Brazil*, 23, 197
- Duncan, R. C., & Thompson, C. 1992, *ApJ*, 392, L9
- Duncan, R. C. 2001, *AIP Conf. Proc.* 586: 20th Texas Symposium on relativistic astrophysics, 586, 495
- Eichler, D. 2002, *MNRAS*, 335, 883
- Figer, D. F., Najarro, F., & Kudritzki, R. P. 2004, *ApJ*, 610, L109
- Fox, D. B., et al. 2005, *Nature*, submitted
- Gavazzi, G., Boselli, A., & Kennicutt, R. 1991, *AJ*, 101, 1207
- George, I. M., & Fabian, A. C. 1991, *MNRAS*, 249, 352
- Ghirlanda, G., Ghisellini, G., & Celotti, A. 2004, *A&A*, 422, L55
- Golenetskii, S. et al., 2004, *GCN circ.* 2923
- Guetta, D. & Piran, T., 2004, Submitted to *A&A*, astro-ph/0407429
- Henden, A. 2002a, *GCN circ.* 1423
- Henden, A. 2002b, *GCN circ.* 1723
- Hurley, K., et al. 2002a, *ApJ*, 567, 447
- Hurley, K., et al. 2002b, *GRB Circular Network*, 1409, 1
- Hurley, K., et al. 2002c, *GRB Circular Network*, 1412, 1
- Hurley, K., et al. 2002d, *GRB Circular Network*, 1719, 1
- Hurley, K., et al. 2003, *GRB Circular Network*, 2492, 1
- Hurley, K., et al. 2005, *Nature*, 434, 1098
- Ibrahim, A. I., Swank, J. H., & Parke, W. 2003, *ApJ*, 584, L17
- Imamura, J. N., & Epstein, R. I. 1987, *ApJ*, 313, 711
- Katz, J. I. 1982, *ApJ*, 260, 371

- Katz, J. I. 1994, *ApJ*, 422, 248
- Katz, J. I., & Canel, L. M. 1996, *ApJ*, 471, 915
- Kennicutt, R. C. 1998, *ARA&A*, 36, 189
- Kouveliotou, C., Meegan, C. A., Fishman, G. J., Bhat, N. P., Briggs, M. S., Koshut, T. M., Paciesas, W. S., & Pendleton, G. N. 1993, *ApJ*, 413, L101
- Kouveliotou, C., et al. 1998, *Nature*, 393, 235
- Lightman, A. P., & White, T. R. 1988, *ApJ*, 335, 57
- Magliocchetti, M., Ghirlanda, G., & Celotti, A. 2003, *MNRAS*, 343, 255
- Mazets, E. P., Cline, T. L., Aptekar, R. L., Frederiks, D. D., Golenetskii, S. V., Il’inskii, V. N., & Pal’shin, V. D. 2005, *ArXiv Astrophysics e-prints*, arXiv:astro-ph/0502541
- Miller, G. E., & Scalo, J. M. 1979, *ApJS*, 41, 513
- Monet, D. G., et al. 2003, *AJ*, 125, 984
- Nakar, E., & Piran, T. 2002, *MNRAS*, 330, 920
- Narayan, R., Paczyński, B., & Piran, T. 1992, *ApJ*, 395, L83
- Paczynski, B. 1992, *Acta Astronomica*, 42, 145
- Palmer, D. M., et al. 2005, *Nature*, 434, 1107
- Pérez-González, P. G., Zamorano, J., Gallego, J., Aragón-Salamanca, A., & Gil de Paz, A. 2003, *ApJ*, 591, 827
- Piran, T., 2004, *Rev. Mod. Phys.*, 76, 1143
- Sakamoto, T., et al. 2005, *GRB Circular Network*, 3010, 1
- Schiminovich et al., 2004, accepted to *ApjL*, astro-ph/0411424
- Schmidt, M. 2001, *ApJ*, 559, L79
- Smith, J. A., et al. 2002, *AJ*, 123, 2121
- Thompson, C., & Duncan, R. C. 1995, *MNRAS*, 275, 255
- Tammann, G. A. 1970, *A&A*, 8, 458

- Timmes, F. X., Diehl, R., & Hartmann, D. H. 1997, *ApJ*, 479, 760
- van den Bergh, S., & Tammann, G. A. 1991, *ARA&A*, 29, 363
- Woods, P. M., Kouveliotou, C., Göğüş, E., Finger, M. H., Swank, J., Markwardt, C. B., Hurley, K., & van der Klis, M. 2002, *ApJ*, 576, 381
- Wyder, T. K., et al. 2005, *ApJ*, 619, L15
- Zacharias, N., Monet, D. G., Levine, S. E., Urban, S. E., Gaume, R., & Wycoff, G. L. 2004, *American Astronomical Society Meeting Abstracts*, 205,

Table 1: The brightest galaxies within short-hard GRB error boxes

GRB	XDSS Plates	RA (J2000)	Dec	NUV [mag]	B [mag]	B-V [mag]	U-V [mag]
031214 ^a	R,I	16:21:44.38	-11:43:07.3	-	18.09±0.3	-	-
021201 ^b	B,R,I	08:07:44.33	21:14:30.1	-	18.88±0.05	1.52±0.05	3.1±0.11
020603 ^c	R,I	15:46:29.36	-22:17:09.9	-	16.03±0.006	1.056±0.005	1.68±0.01
		15:46:34.50	-22:14:52.9	-	16.524±0.002	0.625±0.004	0.71±0.01
001204	B,R,I	02:41:04.36	12:52:39.2	21.3±0.2	19.87 ± 0.07	0.46 ± 0.1	0.34 ± 0.2
000607 ^d	B,R,I	02:33:48.50	17:03:59.5	22.2±0.5	18.15 ± 0.07	0.4 ± 0.1	0.2 ± 0.2
		02:33:47.62	17:03:28.9	22.1±0.4	-	-	-
790613	-	14:11:59.93	78:41:07.0	21.4±0.2	-	-	-

a) This is the magnitude of the brightest clearly resolved galaxy. However, on this plate it was difficult to distinguish nearly-saturated stars from compact round galaxies. We therefore conservatively adopt the brightest nearly-saturated/compact object as our “brightest galaxy” ($B \sim 17$) and assign it (again, conservatively) a very blue color ($U - V = 0$ mag).

b) Both the brightest galaxy and the only other detectable galaxy within the error box, have very red cataloged colors (yielding values of $U - V > 2.5$), and are thus expected to have negligible SFR. We have therefore conservatively assumed that the putative host is blue ($U - V = 0$) and it lies just below the blue plate limit ($B \sim 20.5$).

c) The brightest galaxy is rather red ($U - V \approx 1.7$ mag) and therefore it is not the galaxy with the largest SFR in the field. The properties of the brightest blue galaxy in the field are listed as well.

d) We have observed the only optically detectable galaxy in this error box using the double spectrograph mounted on the 200” telescope at Palomar Observatory. We determine the redshift of this galaxy to be $z = 0.14$, based on detected emission lines (H_α and NII). The lower bound on the distance of a putative host galaxy is derived from a second galaxy with comparable NUV brightness.

Table 2: The minimal distances to putative star-forming host galaxies and corresponding energetics

GRB	$S_{0.015-5\text{MeV}}^a$	$S_{50-300\text{keV}}^a$	D_{min}^b	$E_{min,0.015-5\text{MeV}}^c$	$E_{min,50-300\text{keV}}^c$	Ref.
031214	3×10^{-4}	$4 \times 10^{-5\dagger}$	80 [20]	20 [1.4]	2 [0.14]	Hurley et al. 2003
021201	-	$6 \times 10^{-7\dagger\dagger}$	400 [100]	-	1 [0.06]	Hurley et al. 2002d
020603	-	$2 \times 10^{-5\dagger\dagger}$	100 [25]	-	2 [0.14]	Hurley et al. 2002b,c
001204	2×10^{-6}	5×10^{-7}	400 [100]	4 [0.2]	1 [0.06]	Hurley et al. 2002a
000607	5×10^{-6}	5×10^{-7}	480 [120]	12 [0.8]	1.2 [0.08]	Hurley et al. 2002a
790613	3×10^{-6}	—	400 [100]	6 [0.4]	-	Barat et al. 1984

a) The fluence in erg/cm^2

b) The minimal distance to the burst in Mpc, at $1[2]\sigma$ confidence level, assuming that the rate of short-hard GRBs follow the SFR.

c) The minimal isotropic energy output of the bursts in units of 10^{49} erg at $1[2]\sigma$ confidence level.

$\dagger$) Calculated from the fluence in 15-5000keV assuming a low frequency photon index $\alpha = -1$ and taking a peak frequency $E_p = 2000$ (Hurley et al. 2003).

$\dagger\dagger$) Calculated from the fluence in 25-100 keV assuming a low frequency photon index $\alpha = -1$ and a peak frequency $E_p > 300$




Fig. 1.— The error box of GRB 000607 superposed on a blue POSS II plate. The brightest galaxy turns out to be at $z=0.14$. All other sources within the error box have spatial light distributions consistent with those of stars.